

# User Manual

KC/GRO-alpha/CXCL1 (KC/GRO-alpha/CXCL1) (Murine) Cat. No. MECXP-01013





#### **Description:**

KC/GRO alpha/CXCL1 coded by CXCL1 gene at chromosome 5 is approximately 63% identity to that of mouse MIP2. KC is also approximately 60% identical to the human GROs. Mouse KC is a potent neutrophil attractant and activator. The functional receptor for KC has been identified as CXCR2. Based on the pattern of KC expression in a number of inflammatory disease models, KC appears to have an important role in inflammation. KC was found to be involved in monocyte arrest on atherosclerotic endothelium and may also play a pathophysiological role in Alzheimer's disease.

#### Source:

Escherichia coli

#### Unit:

1 mg

## **Reconstitution:**

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at < -20°C. Further dilutions should be made in appropriate buffered solutions.

## **Formulation:**

Lyophilized from a 0.2  $\mu$ m filtered concentrated solution in PBS, pH 7.4.

#### Storage:

This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.

## **Molecular Weight:**



Approximately 7.8 kDa, a single non-glycosylated polypeptide chain containing 72 amino acid residues.

# **Endotoxin:**

Less than 1 EU/ $\mu$ g of KC/GRO-alpha/CXCL1 as determined by LAL method.

# Usage:

This material is offered by Cyagen Biosciences for research, laboratory or further evaluation purposes. FOR RESEARCH USE ONLY. NOT INTENDED FOR ANY ANIMAL OR HUMAN THERAP EUTIC OR DIAGNOSTIC USE.

## **Biological Activity:**

The biologically active determined by a chemotaxis bioassay using human peripheral blood neutrophils is in a concentration range of 10-100 ng/mL.

## Physical Appearance:

Sterile filtered white lyophilized (freeze-dried) powder.

## AA Sequence:

APIANELRCQ CLQTMAGIHL KNIQSLKVLP SGPHCTQTEV IATLKNGREA CLDPEAPLVQ KIVQKMLKGV PK

## **Purity:**

> 97% by SDS-PAGE and HPLC analyses.



Material Safety Data Sheets (MSDSs) are available upon request.

The Certificate of Analysis (COA), which provides detailed quality control information for each product, is also available at the Cyagen website.

Cyagen Biosciences reserves all rights on the technical documents of its culture products. No part of this document may be reproduced or adapted for other purposes without written permission from Cyagen Biosciences.