



User Manual

Fibroblast Growth Factor-basic (bFGF) (Human)

Cat. No. HEGFP-06021



Description:

Fibroblast Growth Factor-basic (bFGF) is a single-chain polypeptide growth factor that plays a significant role in the process of wound healing and is a potent inducer of angiogenesis. Several different forms of the human protein exist ranging from 18-24 kDa in size due to the use of alternative start sites within the *fgf-2* gene. It has a 55 percent amino acid residue identity to FGF-1 and has potent heparin-binding activity. The growth factor is an extremely potent inducer of DNA synthesis in a variety of cell types from mesoderm and neuroectoderm lineages. It was originally named basic fibroblast growth factor based upon its chemical properties and to distinguish it from acidic fibroblast growth factor. Other homologous FGF belonging to the same family are int-2 (FGF-3), FGF-5 , FGF-6 , K-FGF and KGF (keratinocyte growth factor = FGF-7). All factors are products of different genes, some of which are Oncogene products (FGF-3, FGF-4, FGF-5).

Source:

Escherichia coli

Unit:

10 µg

Reconstitution:

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at < -20°C. Further dilutions should be made in appropriate buffered solutions.

Formulation:

Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4.

Storage:

This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.

Molecular Weight:

Approximately 16.5 kDa, a single non-glycosylated polypeptide chain containing 147 amino acids.

Endotoxin:

Less than 1 EU/μg of bFGF as determined by LAL method.

Usage:

This material is offered by Cyagen Biosciences for research, laboratory or further evaluation purposes. FOR RESEARCH USE ONLY. NOT INTENDED FOR ANY ANIMAL OR HUMAN THERAP EUTIC OR DIAGNOSTIC USE.

Biological Activity:

The ED50 determined by a cell proliferation assay using murine balb/c 3T3 cells is less than 0.6 ng/mL, corresponding to a specific activity of $> 1.7 \times 10^6$ IU/mg.

Physical Appearance:

Sterile filtered white lyophilized (freeze-dried) powder.

AA Sequence:

MPALPEDGGS GAFPPGHFKD PKRLYCKNGG FFLRIHPDGR VDGVREKSDP
HIKLQLQAE E RGVVSIKVC ANRYLAMKED GRLLASKCVT DECFFFERLE
SNNYNTYRSR KYTSWYVALK RTGQYKLGSK TGPGQKAILF LPMSAKS

Purity:

> 96% by SDS-PAGE and HPLC analyses.

Material Safety Data Sheets (MSDSs) are available upon request.

The Certificate of Analysis (COA), which provides detailed quality control information for each product, is also available at the Cyagen website.

Cyagen Biosciences reserves all rights on the technical documents of its culture products. No part of this document may be reproduced or adapted for other purposes without written permission from Cyagen Biosciences.