



## User Manual

**Granulocyte Colony Stimulating Factor (G-CSF) (Human)**  
**Cat. No. HEOPP-07013**



**Description:**

Granulocyte Colony Stimulating Factor (G-CSF) is a pleiotropic cytokine best known for its specific effects on the proliferation, differentiation, and activation of hematopoietic cells of the neutrophilic granulocyte lineage. It is produced mainly by monocytes and macrophages upon activation by endotoxin, TNF- $\alpha$  and IFN- $\gamma$ . Other cell types including fibroblasts, endothelial cells, astrocytes and bone marrow stromal cells can also secrete G-CSF after LPS, IL-1 or TNF- $\alpha$  activation. In addition, various carcinoma cell lines and myeloblastic leukemia cells can express G-CSF constitutively. In humans, two distinct cDNA clones for G-CSF, encoding 207 and 204 amino acid precursor proteins, have been isolated. Both proteins have a 30 amino acid signal peptide and have identical amino acid sequences except for a three amino acid insertion (deletion) at the 35th amino acid residue from the N-terminus of the mature protein. Human G-CSF is 73% identical at the amino acid level to murine G-CSF and the two proteins show species cross-reactivity.

**Source:**

*Escherichia coli*

**Unit:**

1 mg

**Reconstitution:**

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at  $< -20^{\circ}\text{C}$ . Further dilutions should be made in appropriate buffered solutions.

**Formulation:**

Lyophilized from a 0.2  $\mu\text{m}$  filtered concentrated solution in 10 mM sodium acetate buffer, containing 5% trehalose, pH 4.0.

**Storage:**

This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.

**Molecular Weight:**

Approximately 18.7 kDa, a single non-glycosylated polypeptide chain containing 174 amino acids.

**Endotoxin:**

Less than 1 EU/μg of G-CSF as determined by LAL method.

**Usage:**

This material is offered by Cyagen Biosciences for research, laboratory or further evaluation purposes. FOR RESEARCH USE ONLY. NOT INTENDED FOR ANY ANIMAL OR HUMAN THERAP EUTIC OR DIAGNOSTIC USE.

**Biological Activity:**

The ED50 determined by a cell proliferation assay using murine NFS-60 cells is less than 0.1 ng/mL, corresponding to a specific activity of  $> 1.0 \times 10^7$  IU/mg.

**Physical Appearance:**

Sterile filtered white lyophilized (freeze-dried) powder.

### **AA Sequence:**

TPLGPASSLP QSFLLKCLEQ VRKIQGDGAA LQEKLCATYK LCHPEELVLL  
GHSLGIPWAP LSSCPSQALQ LAGCLSQLHS GLFLYQGLLQ ALEGISPELG  
PTLDTLQLDV ADFATTIWQQ MEELGMAPAL QPTQGAMPAF ASAFQRRAGG  
VLVASHLQSF LEVSYRVLRH LAQP

### **Purity:**

> 98% by SDS-PAGE and HPLC analyses.

Material Safety Data Sheets (MSDSs) are available upon request.

The Certificate of Analysis (COA), which provides detailed quality control information for each product, is also available at the Cyagen website.

Cyagen Biosciences reserves all rights on the technical documents of its culture products. No part of this document may be reproduced or adapted for other purposes without written permission from Cyagen Biosciences.