



## User Manual

**Bone Morphogenetic Protein-7 (BMP-7) (Human)**

**Cat. No. HEOPP-02103**



**Description:**

Human Bone Morphogenetic Protein-7 (BMP-7) is one of at least 15 structurally and functionally related BMPs, which are members of the transforming growth factor  $\beta$  (TGF- $\beta$ ) superfamily. BMPs were originally identified as protein regulators of cartilage and bone formation. However, they have since been shown to be involved in embryogenesis and morphogenesis of various tissues and organs. BMPs have also been shown to regulate the growth, differentiation, chemotaxis and apoptosis of various cell types, including mesenchymal cells, epithelial cells, hematopoietic cells and neuronal cells. BMP-7 is synthesized as large precursor molecules which are cleaved by proteolytic enzymes. The active form can consist of a dimer of two identical proteins or a heterodimer of two related bone morphogenetic proteins.

**Source:**

*Escherichia coli*

**Unit:**

1 mg

**Reconstitution:**

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in 10 mM HAc to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at  $< -20^{\circ}\text{C}$ . Further dilutions should be made in appropriate buffered solutions.

**Formulation:**

Lyophilized from a 0.2  $\mu\text{m}$  filtered concentrated solution in 30% acetonitrile, 0.1% TFA.

**Storage:**

This lyophilized preparation is stable at 2-8 $^{\circ}\text{C}$ , but should be kept at -20 $^{\circ}\text{C}$  for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8 $^{\circ}\text{C}$ . For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20 $^{\circ}\text{C}$  to -70 $^{\circ}\text{C}$ . Avoid repeated freeze/thaw cycles.

**Molecular Weight:**

Approximately 15.6 kDa, a monomeric, non-glycosylated polypeptide chain containing 139 amino acids.

**Endotoxin:**

Less than 1 EU/μg of BMP-7 as determined by LAL method.

**Usage:**

This material is offered by Cyagen Biosciences for research, laboratory or further evaluation purposes. FOR RESEARCH USE ONLY. NOT INTENDED FOR ANY ANIMAL OR HUMAN THERAP EUTIC OR DIAGNOSTIC USE.

**Biological Activity:**

The ED50 determined by inducing alkaline phosphatase production of murine ATDC5 cells is less than 600 ng/mL, correspond-ing to a specific activity of  $> 1.7 \times 10^3$  IU/mg.

**Physical Appearance:**

Sterile filtered white lyophilized (freeze-dried) powder.

**AA Sequence:**

STGSKQRSQN RSKTPKNQEA LRMANVAENS SSDQRQACKK HELYVSFRDL  
GWQDWIIAPE GYAAYYCEGE CAFPLNSYMN ATNHAIVQTL VHFINPETVP  
KPCCAPTQLN AISVLYFDDS SNVILKKYRN MVVRACGCH

**Application:**

1. Molecular standard (Western, ELISA) in studying secreted BMP-7; 2. Preparing antibodies for BMP-7 monomer; 3. Molecule standard in detecting secreted BMP-7 in reduced SDS-PAGE.

**Purity:**

> 95% by SDS-PAGE and HPLC analyses.

Material Safety Data Sheets (MSDSs) are available upon request.

The Certificate of Analysis (COA), which provides detailed quality control information for each product, is also available at the Cyagen website.

Cyagen Biosciences reserves all rights on the technical documents of its culture products. No part of this document may be reproduced or adapted for other purposes without written permission from Cyagen Biosciences.