



User Manual

NAP-2/CXCL7 (NAP-2/CXCL7) (Human)

Cat. No. HECXP-07013

Description:

Neutrophil Activating Peptide 2 (NAP-2) is proteolytically processed carboxyl-terminal fragments of platelet basic protein (PBP) which is found in the alpha-granules of human platelets. NAP-2 is a member of the CXC chemokines. Similar to other ELR domain containing CXC chemokines such as IL-8 and the GRO proteins, NAP-2 has been shown to bind CXCR-2 and to chemoattract and activate neutrophils. Although CTAP-III, β -TG and PBP represent amino-terminal extended variants of NAP-2 and possess the same CXC chemokine domains, these proteins do not exhibit NAP-2 activity. Recently, it has been shown that the additional amino-terminal residues of CTAP-III masks the critical ELR receptor binding domain that is exposed on NAP-2 and may account for lack of NAP-2 activity.

Source:

Escherichia coli

Unit:

1 mg

Reconstitution:

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at $< -20^{\circ}\text{C}$. Further dilutions should be made in appropriate buffered solutions.

Formulation:

Lyophilized from a 0.2 μm filtered concentrated solution in 20 mM PB, pH 7.4, 50 mM NaCl.

Storage:

This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.

Molecular Weight:

Approximately 7.6 kDa, a single non-glycosylated polypeptide chain containing 70 amino acids.

Endotoxin:

Less than 1 EU/μg of NAP-2/CXCL7 as determined by LAL method.

Usage:

This material is offered by Cyagen Biosciences for research, laboratory or further evaluation purposes. FOR RESEARCH USE ONLY. NOT INTENDED FOR ANY ANIMAL OR HUMAN THERAP EUTIC OR DIAGNOSTIC USE.

Biological Activity:

The biologically active determined by a chemotaxis bioassay using human peripheral blood neutrophils is in a concentration range of 0.1-1.0 ng/mL.

Physical Appearance:

Sterile filtered white lyophilized (freeze-dried) powder.

AA Sequence:

AELRCMCIKT TSGIHPKNIQ SLEVIGKGTH CNQVEVIATL KDGRKICLDP DAPRIKKIVQ
KKLAGDESAD

Purity:

> 97% by SDS-PAGE and HPLC analyses.

Material Safety Data Sheets (MSDSs) are available upon request.

The Certificate of Analysis (COA), which provides detailed quality control information for each product, is also available at the Cyagen website.

Cyagen Biosciences reserves all rights on the technical documents of its culture products. No part of this document may be reproduced or adapted for other purposes without written permission from Cyagen Biosciences.