



User Manual

Beta Defensin-3 (BD-3) (Murine)

Cat. No. MEOPP-02033

Description:

Beta defensin-3, also known as BD-3 and DEFB-3, is a membrane active cationic peptide that functions in inflammation and innate immune responses and coded by Defb 3 gene on chromosome 8 in mouse. There are at least 30 β -defensins which are distinguished from α -defensins by the connectivity pattern of their three intramolecular disulfide bonds. BD3 is widely expressed among epithelial tissues, notably by keratinocytes and airway epithelial cells. It is upregulated in response to proinflammatory cytokines, microbial and viral infections, and at the edges of skin wounds. BD3 induction in osteoarthritis chondrocytes promotes MMP1 and 13 productions and inhibits TIMP1 and 2 expressions.

Source:

Escherichia coli

Unit:

1 mg

Reconstitution:

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at $< -20^{\circ}\text{C}$. Further dilutions should be made in appropriate buffered solutions.

Formulation:

Lyophilized from a 0.2 μm filtered concentrated solution in 2 \times PBS, pH 7.4.

Storage:

This lyophilized preparation is stable at 2-8 $^{\circ}\text{C}$, but should be kept at -20 $^{\circ}\text{C}$ for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one

week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.

Molecular Weight:

Approximately 4.6 kDa, a single non-glycosylated polypeptide chain containing 41 amino acid residues.

Endotoxin:

Less than 1 EU/μg of BD-3 as determined by LAL method.

Usage:

This material is offered by Cyagen Biosciences for research, laboratory or further evaluation purposes. FOR RESEARCH USE ONLY. NOT INTENDED FOR ANY ANIMAL OR HUMAN THERAP EUTIC OR DIAGNOSTIC USE.

Biological Activity:

The ED50 determined by anti-microbial activity against E.coli. is less than 20 μg/mL, corresponding to a specific activity of > 50 IU/mg.

Physical Appearance:

Sterile filtered white lyophilized (freeze-dried) powder.

AA Sequence:

KKINNPVSCL RKGGRWCNRC IGNTRQIGSC GVPFLKCKR K

Purity:

> 95% by SDS-PAGE and HPLC analyses.

Material Safety Data Sheets (MSDSs) are available upon request.

The Certificate of Analysis (COA), which provides detailed quality control information for each product, is also available at the Cyagen website.

Cyagen Biosciences reserves all rights on the technical documents of its culture products. No part of this document may be reproduced or adapted for other purposes without written permission from Cyagen Biosciences.