



User Manual

Beta Defensins-2 (BD-2) (Murine)

Cat. No. MEOPP-02022



Description:

Defensins (alpha and beta) are cationic peptides with a broad spectrum of antimicrobial activity that comprise an important arm of the innate immune system. The α -defensins are distinguished from the β -defensins by the pairing of their three disulfide bonds. To date, four β -defensins have been identified: BD-1, BD-2, BD-3 and BD-4. β -defensins are expressed on some leukocytes and at epithelial surfaces. In addition to their direct antimicrobial activities, they are chemoattractant towards immature dendritic cells and memory T cells. The β -defensin proteins are expressed as the C-terminal portion of precursors and are released by proteolytic cleavage of a signal sequence and, in the case of BD-1 (36 a.a.), a propeptide region. β -defensins contain a six-cysteine motif that forms three intra-molecular disulfide bonds.

Source:

Escherichia coli

Unit:

20 μ g

Reconstitution:

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at $< -20^{\circ}\text{C}$. Further dilutions should be made in appropriate buffered solutions.

Formulation:

Lyophilized from a 0.2 μm filtered concentrated solution in PBS, pH 7.4.

Storage:

This lyophilized preparation is stable at 2-8 $^{\circ}\text{C}$, but should be kept at -20 $^{\circ}\text{C}$ for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8 $^{\circ}\text{C}$. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20 $^{\circ}\text{C}$ to -70 $^{\circ}\text{C}$. Avoid repeated freeze/thaw cycles.

Molecular Weight:

Approximately 5.54 kDa, a single non-glycosylated polypeptide chain containing 51 amino acids.

Endotoxin:

Less than 1 EU/ μ g of BD-2 as determined by LAL method.

Usage:

This material is offered by Cyagen Biosciences for research, laboratory or further evaluation purposes. FOR RESEARCH USE ONLY. NOT INTENDED FOR ANY ANIMAL OR HUMAN THERAPEUTIC OR DIAGNOSTIC USE.

Biological Activity:

The biological activity determined by a chemotaxis bioassay using immature human dendritic cells is in a concentration of 10-100 ng/mL.

Physical Appearance:

Sterile filtered white lyophilized (freeze-dried) powder.

AA Sequence:

AVGSLKSIQY EAELDHCHT N GGYCVRAICP PSARRPGSCF PEKNPCCKYM K

Purity:

> 98% by SDS-PAGE and HPLC analyses.

Material Safety Data Sheets (MSDSs) are available upon request.

The Certificate of Analysis (COA), which provides detailed quality control information for each product, is also available at the Cyagen website.

Cyagen Biosciences reserves all rights on the technical documents of its culture products. No part of this document may be reproduced or adapted for other purposes without written permission from Cyagen Biosciences.