



User Manual

Tumor Necrosis Factor-alpha (TNF- α) (Rat)

Cat. No. RETNP-01012

Description:

Tumor necrosis factor-alpha (TNF- α), also called cachectin, is produced by neutrophils, activated lymphocytes, macrophages, NK cells, LAK cells, astrocytes endothelial cells, smooth muscle cells and some transformed cells. TNF- α occurs as a secreted, soluble form and as a membrane-anchored form, both of which are biologically active. The naturally-occurring form of TNF- α is glycosylated, but non-glycosylated recombinant TNF- α has comparable biological activity. The biologically active native form of TNF- α is reportedly a trimer. Two types of receptors for TNF- α have been described and virtually all cell types studied show the presence of one or both of these receptor types.

Source:

Escherichia coli

Unit:

20 μ g

Reconstitution:

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at $< -20^{\circ}\text{C}$. Further dilutions should be made in appropriate buffered solutions.

Formulation:

Lyophilized from a 0.2 μ m filtered concentrated solution in 20 mM PB, pH7.2, 150 mM NaCl.

Storage:

This lyophilized preparation is stable at 2-8 $^{\circ}\text{C}$, but should be kept at -20 $^{\circ}\text{C}$ for long term

storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.

Molecular Weight:

Approximately 17.2 kDa, a single non-glycosylated polypeptide chain containing 156 amino acids.

Endotoxin:

Less than 1 EU/μg of TNF-α as determined by LAL method.

Usage:

This material is offered by Cyagen Biosciences for research, laboratory or further evaluation purposes. FOR RESEARCH USE ONLY. NOT INTENDED FOR ANY ANIMAL OR HUMAN THERAP EUTIC OR DIAGNOSTIC USE.

Biological Activity:

The ED50 determined by a cytotoxicity assay using murine L929 cells is less than 0.1 ng/mL, corresponding to a specific activity of $> 1.0 \times 10^7$ IU/mg in the presence of actinomycin D.

Physical Appearance:

Sterile filtered white lyophilized (freeze-dried) powder.

AA Sequence:

LRSSSQNSSD KPVAHVVANH QAEEQLEWLS QRANALLANG MDLKDNQLVV
PADGLYLIYS QVLFKGGQCP DYVLLTHTVS RFAISYQEKV SLLSAIKSPC

PKDTPEGAEL KPWYEPMYLG GVFQLEKGD LSAEVLNPKY LDITESGQVY FGVIAL

Purity:

> 98% by SDS-PAGE and HPLC analyses.

Material Safety Data Sheets (MSDSs) are available upon request.

The Certificate of Analysis (COA), which provides detailed quality control information for each product, is also available at the Cyagen website.

Cyagen Biosciences reserves all rights on the technical documents of its culture products. No part of this document may be reproduced or adapted for other purposes without written permission from Cyagen Biosciences.