



User Manual

SDF-1 alpha/CXCL12 (SDF-1 α /CXCL12) (Human)

Cat. No. HECXP-12013

Description:

SDF-1 α and SDF-1 β , members of the chemokine α subfamily that lack the ELR domain, were initially identified using the signal sequence trap cloning strategy from a mouse bone-marrow stromal cell line. These proteins were subsequently also cloned from a human stromal cell line as cytokines that supported the proliferation of a stromal cell-dependent pre-B-cell line. SDF-1 α and SDF-1 β cDNAs encode precursor proteins of 89 and 93 amino acid residues, respectively. Both SDF-1 α and SDF-1 β are encoded by a single gene and arise by alternative splicing. The two proteins are identical except for the four amino acid residues that are present in the carboxy-terminus of SDF-1 β and absent from SDF-1 α . SDF-1/PBSF is highly conserved between species, with only one amino acid substitution between the mature human and mouse proteins. SDF-1/PBSF acts via the chemokine receptor CXCR4 and has been shown to be a chemoattractant for T-lymphocytes, monocytes, pro- and pre- B cells, but not neutrophils.

Source:

Escherichia coli

Unit:

1 mg

Reconstitution:

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at $< -20^{\circ}\text{C}$. Further dilutions should be made in appropriate buffered solutions.

Formulation:

Lyophilized from a 0.2 μm filtered concentrated solution in 20 mM PB, pH 7.0, 130 mM NaCl.

Storage:

This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.

Molecular Weight:

Approximately 8.0 kDa, a single non-glycosylated polypeptide chain containing 68 amino acids.

Endotoxin:

Less than 1 EU/μg of SDF-1α/CXCL12 as determined by LAL method.

Usage:

This material is offered by Cyagen Biosciences for research, laboratory or further evaluation purposes. FOR RESEARCH USE ONLY. NOT INTENDED FOR ANY ANIMAL OR HUMAN THERAP EUTIC OR DIAGNOSTIC USE.

Biological Activity:

The biologically active determined by a chemotaxis bioassay using PHA and IL-2 activated human peripheral blood T-lymphocytes is in a concentration range of 20-80 ng/mL.

Physical Appearance:

Sterile filtered white lyophilized (freeze-dried) powder.

AA Sequence:

KPVLSYRCP CRFFESHVAR ANVKHLKILN TPNCALQIVA RLKNNNRQVC
IDPKLKWIQE YLEKALNK

Purity:

> 97% by SDS-PAGE and HPLC analyses.

Material Safety Data Sheets (MSDSs) are available upon request.

The Certificate of Analysis (COA), which provides detailed quality control information for each product, is also available at the Cyagen website.

Cyagen Biosciences reserves all rights on the technical documents of its culture products. No part of this document may be reproduced or adapted for other purposes without written permission from Cyagen Biosciences.