



## User Manual

**Eotaxin-3/CCL26 (Eotaxin-3/CCL26) (Human)**

**Cat. No. HECCP-26012**

**Description:**

Eotaxin3, also named CCL26 or SCYA26, is a novel human CC chemokine coded by CXCL26 gene at chromosome 7 in human. Recombinant Eotaxin3/CCL26 has been produced in insect cells using a baculovirus expression system and shown to contain 71 aa residues. Recombinant Eotaxin3/CCL26 is chemotactic for eosinophils and PHAactivated T cells. Eotaxin3/CCL26 induces calcium flux in eosinophils as well as in CCR3 transfected cells. Eotaxin3/CCL26 has also been shown to crossdesensitize cells to other CCR3 ligands. Both the 71 aa residue and 68 aa residue variants of recombinant Eotaxin3 have been expressed in *E. coli* and found to have equal potency in inducing chemotaxis of a human CCR3 transfected cell line.

**Source:**

*Escherichia coli*

**Unit:**

20 µg

**Reconstitution:**

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at < -20°C. Further dilutions should be made in appropriate buffered solutions.

**Formulation:**

Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4.

**Storage:**

This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term

storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.

### **Molecular Weight:**

Approximately 8.4 kDa, a single non-glycosylated polypeptide chain containing 71 amino acid residues.

### **Endotoxin:**

Less than 1 EU/μg of Eotaxin-3/CCL26 as determined by LAL method.

### **Usage:**

This material is offered by Cyagen Biosciences for research, laboratory or further evaluation purposes. FOR RESEARCH USE ONLY. NOT INTENDED FOR ANY ANIMAL OR HUMAN THERAP EUTIC OR DIAGNOSTIC USE.

### **Biological Activity:**

The biologically active determined by a chemotaxis bioassay using murine CCR3 transfected murine BaF3 cells is in a concentration range of 0.5-1.5 μg/mL.

### **Physical Appearance:**

Sterile filtered white lyophilized (freeze-dried) powder.

### **AA Sequence:**

TRGSDISKTC CFQYSHKPLP WTWVRSYEFT SNSCSQRAVI FTTKRGKKVC  
THPRKKWVQK YISLLKTPKQ L

**Purity:**

> 97% by SDS-PAGE and HPLC analyses.

Material Safety Data Sheets (MSDSs) are available upon request.

The Certificate of Analysis (COA), which provides detailed quality control information for each product, is also available at the Cyagen website.

Cyagen Biosciences reserves all rights on the technical documents of its culture products. No part of this document may be reproduced or adapted for other purposes without written permission from Cyagen Biosciences.