



## User Manual

**Epidermal Growth Factor (EGF) (Rat)**

**Cat. No. REGFP-05011**

**Description:**

Epidermal Growth Factor EGF is a potent growth factor that stimulates the proliferation of various epidermal and epithelial cells. Additionally, EGF has been shown to inhibit gastric secretion, and to be involved in wound healing. EGF signals through a receptor known as c-erbB, which is a class I tyrosine kinase receptor. This receptor also binds with TGF- $\alpha$  and VGF (vaccinia virus growth factor).

**Source:**

*Escherichia coli*

**Unit:**

20  $\mu$ g

**Reconstitution:**

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at  $< -20^{\circ}\text{C}$ . Further dilutions should be made in appropriate buffered solutions.

**Formulation:**

Lyophilized from a 0.2  $\mu$ m filtered solution in PBS, pH 7.4.

**Storage:**

This lyophilized preparation is stable at 2-8 $^{\circ}\text{C}$ , but should be kept at -20 $^{\circ}\text{C}$  for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8 $^{\circ}\text{C}$ . For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20 $^{\circ}\text{C}$  to -70 $^{\circ}\text{C}$ . Avoid repeated freeze/thaw cycles.

**Molecular Weight:**

Approximately 6.2 kDa, a single non-glycosylated polypeptide chain containing 53 amino acids, including 3 intramolecular disulfide-bonds.

**Endotoxin:**

Less than 1 EU/μg of EGF as determined by LAL method.

**Usage:**

This material is offered by Cyagen Biosciences for research, laboratory or further evaluation purposes. FOR RESEARCH USE ONLY. NOT INTENDED FOR ANY ANIMAL OR HUMAN THERAP EUTIC OR DIAGNOSTIC USE.

**Biological Activity:**

The ED50 determined by a cell proliferation assay using murine Balb/c 3T3 cells is less than 0.1 ng/mL, corresponding to a specific activity of  $> 1.0 \times 10^7$  IU/mg.

**Physical Appearance:**

Sterile filtered white lyophilized (freeze-dried) powder.

**AA Sequence:**

NSNTGCPPSY DGYCLNGGVC MYVESVDRYV CNCVIGYIGE RCQHRDLRWW KLR

**Purity:**

97% by SDS-PAGE and HPLC analyses.

Material Safety Data Sheets (MSDSs) are available upon request.

The Certificate of Analysis (COA), which provides detailed quality control information for each product, is also available at the Cyagen website.

Cyagen Biosciences reserves all rights on the technical documents of its culture products. No part of this document may be reproduced or adapted for other purposes without written permission from Cyagen Biosciences.