



User Manual

ENA-78/CXCL5 (8-78a.a.) (ENA-78/CXCL5 (8-78a.a.)) (Human)
Cat. No. HECXP-05023

Description:

Epithelial cell-derived neutrophil-activating peptide 78 (ENA-78) is a member of the CXC subfamily of chemokines that has the Glu-Leu-Arg (ELR) motif preceding the CXC motif. Similar to other ELR containing CXC chemokines, ENA-78 is a potent neutrophil chemoattractant and activator. Proteolysis of ENA-78 with cathepsin G and chymotrypsin have yielded N-terminally truncated variants with increased biological activities. ENA-70 and ENA-74 represent truncated recombinant ENA-78 variants missing 8 and 4 aa residues, respectively, from the N-terminus. Recombinant ENA-70 and ENA-74 have been shown to have increased potency in neutrophil chemotaxis and myeloperoxidase and elastase release assays.

Source:

Escherichia coli

Unit:

1 mg

Reconstitution:

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at < -20°C. Further dilutions should be made in appropriate buffered solutions.

Formulation:

Lyophilized from a 0.2 µm filtered concentrated solution in 2xPBS, pH 7.4.

Storage:

This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term

storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.

Molecular Weight:

Approximately 7.8 kDa, a single non-glycosylated polypeptide chain containing 71 amino acids.

Endotoxin:

Less than 1 EU/μg of ENA-78/CXCL5 (8-78a.a.) as determined by LAL method.

Usage:

This material is offered by Cyagen Biosciences for research, laboratory or further evaluation purposes. FOR RESEARCH USE ONLY. NOT INTENDED FOR ANY ANIMAL OR HUMAN THERAP EUTIC OR DIAGNOSTIC USE.

Biological Activity:

The biologically active determined by a chemotaxis bioassay using human peripheral blood neutrophils is in a concentration of 5.0-10 ng/mL.

Physical Appearance:

Sterile filtered white lyophilized (freeze-dried) powder.

AA Sequence:

LRELRCVCLQ TTQGVHPKMI SNLQVFAIGP QCSKVEVVAS LKNGKEICLD
PEAPFLKKVI QKILDGGNKE N

Purity:

> 95% by SDS-PAGE and HPLC analyses.

Material Safety Data Sheets (MSDSs) are available upon request.

The Certificate of Analysis (COA), which provides detailed quality control information for each product, is also available at the Cyagen website.

Cyagen Biosciences reserves all rights on the technical documents of its culture products. No part of this document may be reproduced or adapted for other purposes without written permission from Cyagen Biosciences.